arXiv:nucl-th/0011084v2 15 Aug 2001

Nucleon and meson effective masses in the Relativistic Mean-Field Theory

Ryszard Mańka ‡ **and Ilona Bednarek** §

Institute of Physics, Silesian University, Uniwersytecka 4, 40-007 Katowice, Poland.

Abstract. Nucleon and meson effective masses in the nonlinear Relativistic Mean-Field Theory (RMF) introducing a nonlinear $\omega - \rho$ and σ coupling motivated by the Quark Meson Coupling model (QMC) is explored. It is shown that, in contrast to the usual Walecka model, not only the effective nucleon mass $m_{eff,N}$ but also the effective σ, ρ meson masses ($m_{eff,\sigma}$, $m_{eff,\rho}$) and the effective ω meson mass $m_{eff,\omega}$ are nucleon density dependent.

PACS numbers: 24.10.Cn, 24.10.Jv, 21.30.Fe

Submitted to: *J. Phys. G: Nucl. Phys.*

1. Introduction

Relativistic mean-field theory seems to be a powerful tool in describing various aspects of nuclear structure [1, 2, 3]. This theory provides an elegant and economical framework, in which properties of nuclear matter [4, 5], finite nuclei and neutron star, as well as the dynamics of heavy-ion collisions, can be calculated. Compared to conventional nonrelativistic approaches, relativistic models explicitly include mesonic degrees of freedom and describe nucleons as Dirac quasiparticles. Moreover, the spin-orbit interaction arises naturally from the Dirac-Lorenz structure of the effective Lagrangian. Nucleons interact in a relativistic covariant manner through the exchange of virtual mesons: the isoscalar scalar σ-meson, the isoscalar vector ω-meson, and the isovector vector ρ-meson. The model is based on the one-boson exchange description of the nucleon-nucleon interaction. One of the most interesting problems in intermediate energy physics which has been investigated with many approaches is an effective meson mass in nuclear medium.

2. Effective vector meson masses in nuclear medium.

In this paper the properties of nuclear matter in high-density and neutron-rich regime are considered with the use of the relativistic mean-field approximation. This approach implies the interaction of nucleons through the exchange of meson fields, so the model considered here comprises: nucleons and scalar, vector-isoscalar and vector-isovector mesons. Consequently contributions coming from these components influence the properties of nucleon. In this paper the nonlinear vector-isoscalar self-interaction is dealt. Modification of this type was proposed by Bodmer in order to achieve good agreement with the Dirac-Brückner [6] calculations at high densities. The first version of the ρ meson field introduction is of a minimal type without any nonlinearity and consists only of the coupling of this field to nucleons. This case is enlarged by the nonlinear vector-isoscalar and vector-isovector interaction and the vector-scalar interactions which modify the density dependence of the ρ mean field and the energy symmetry. Such an extension was inspired by the paper [7] in which the authors indicate the existence of a relation between the neutron-rich skin of a heavy nucleus and the properties of a neutron star crust. In the quark meson coupling model [9] nucleon properties are modified by the meson coupling directly to the quarks and not to the nucleons (paper by Li et.al. [8]). The RMF theory is an effective one, as coupling constants are determined by the bulk properties of nuclear matter such as saturation density, binding energy, compressibility and the symmetry energy.

The starting point in this paper is the construction of the effective Lagrangian [10, 11, 12] intended for application to the system described above. The scalar σ, isoscalar-vector ω and isovector-vector ρ mesons are denoted by φ, ω_μ, b^a_μ, respectively. The Lagrange density function for this model has the following form

$$\mathcal{L} = \frac{1}{2}\partial_\mu\varphi\partial^\mu\varphi - U(\varphi) - \frac{1}{4}\Omega_{\mu\nu}\Omega^{\mu\nu} + \frac{1}{2}(M_\omega - g_{\omega\sigma}\varphi)^2\omega_\mu\omega^\mu + \frac{1}{4}c_3(\omega_\mu\omega^\mu)^2$$

$$-\frac{1}{4}R^{a}_{\mu\nu}R^{a\mu\nu}+\frac{1}{2}(M_\rho-g_{\rho\sigma}\varphi)^2b^a_\mu b^{a\mu}+\Lambda_v(g_\rho g_\omega)^2(\omega_\mu\omega^\mu)(b^a_\mu b^{a\mu})$$
$$-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}+i\overline{\psi}\gamma^\mu D_\mu\psi-\overline{\psi}(M-g_{N\sigma}\varphi)\psi. \qquad (1)$$

The field tensors $R^a_{\mu\nu}$, $\Omega_{\mu\nu}$, $F_{\mu\nu}$ and the covariant derivative D_μ are given by

$$R^a_{\mu\nu}=\partial_\mu b^a_\nu-\partial_\nu b^a_\mu+g_\rho\varepsilon_{abc}b^b_\mu b^c_\nu, \qquad (2)$$

$$\Omega_{\mu\nu}=\partial_\mu\omega_\nu-\partial_\nu\omega_\mu, \qquad (3)$$

$$F_{\mu\nu}=\partial_\mu A_\nu-\partial_\nu A_\mu, \qquad (4)$$

$$D_\mu=\partial_\mu+\frac{1}{2}ig_\rho b^a_\mu\tau^a+ig_\omega\omega_\mu+iQ_eA_\mu. \qquad (5)$$

The potential function $U(\varphi)$ possesses a polynomial form introduced by Boguta and Bodmer [13] in order to get a correct value of the compressibility K of nuclear matter at saturation density

$$U(\varphi)=\frac{1}{2}m_s^2\varphi^2+\frac{1}{3}g_2\varphi^3+\frac{1}{4}g_3\varphi^4. \qquad (6)$$

The nucleon field ψ has a form of a column matrix composed of proton and neutron fields, respectively

$$\psi=\left(\begin{array}{c}\psi_n\\ \psi_p\end{array}\right). \qquad (7)$$

The nucleon mass is denoted by M whereas m_s, M_ω, M_ρ are masses assigned to the meson fields. The parameters entering the Lagrangian function (1) are the coupling constants c_3, g_ω, g_ρ, $g_{N\sigma}$ (Table 3) for meson fields, $e^2/4\pi=1/137$ for photon field and the self-interacting coupling constants g_2 and g_3. The coupling constants $g_{\omega\sigma}$ and $g_{\rho\sigma}$ in the QMC model [14] are related to the constant $g_{N\sigma}$ by the relations

$$g_{\omega\sigma}=\frac{2}{3}g_{N\sigma},\ g_{\rho\sigma}=\frac{2}{3}g_{N\sigma}. \qquad (8)$$

The Lagrangian function (1) includes also nonlinear terms

$$\frac{1}{4}c_3(\omega_\mu\omega^\mu)^2+\Lambda_v(g_\rho g_\omega)^2(\omega_\mu\omega^\mu)(b^a_\mu b^{a\mu}) \qquad (9)$$
$$-g_{\omega\sigma}M_\omega\varphi\,\omega_\mu\omega^\mu+\frac{1}{2}g^2_{\omega\sigma}\varphi^2\omega_\mu\omega^\mu-g_{\rho\sigma}M_\omega\varphi\,b^a_\mu b^{a\mu}+\frac{1}{2}g^2_{\rho\sigma}\varphi^2b^a_\mu b^{a\mu}$$

which affect remarkably properties of nucleons. The first term was proposed by Bodmer in order to achieve good agreement with the Dirac-Brückner [6] calculations at high densities. All terms which involve the b^a_μ field change the density dependence of the symmetry energy and cause the change in the density dependence of the ρ field. The modification of the density dependence of the ρ mean field (5) can lead to the change of the neutron-skin thickness [7]. This effect is observed in the Parity Radius Experiment (PREX) at the Jefferson Laboratory which aims to measure the neutron radius in ^{208}Pb

via parity violating electron scattering [15].
The field equations for meson fields $\Phi_A = \{\varphi,\, \omega_\mu,\, b^a_\mu\}$ are

$$\begin{aligned}\Box\varphi &= m_s^2\varphi + g_2\varphi^2 + g_3\varphi^3 - g_{\sigma N}\overline{\psi}\psi + (M_\omega - g_{\omega\sigma}\varphi)g_{\omega\sigma}\omega_\mu\omega^\mu \\ &\quad + (M_\rho - g_{\rho\sigma}\varphi)g_{\rho\sigma}b^a_\mu b^{a\mu},\end{aligned} \tag{10}$$

$$\begin{aligned}-\partial_\mu\Omega^{\mu\nu} &= (M_\omega - g_{\omega\sigma}\varphi)^2\omega^\nu + c_3(\omega_\mu\omega^\mu)\omega^\nu \\ &\quad + 2\Lambda_v(g_\omega g_\rho)^2(b^a_\mu b^{a\mu})\omega^\nu - g_\omega J^\nu_B,\end{aligned} \tag{11}$$

$$-D_\mu R^{\mu\nu a} = (M_\rho - g_{\rho\sigma}\varphi)^2 b^{\nu a} + 2\Lambda_v(g_\omega g_\rho)^2(\omega_\mu\omega^\mu)b^{a\nu} - g_\rho\delta_{a,3}J^\nu_3. \tag{12}$$

Sources that appear in these equations are the baryon current

$$J^\nu_B = \overline{\psi}\gamma^\nu\psi \tag{13}$$

and the isospin current which exists only in the asymmetric matter

$$J^\nu_3 = \frac{1}{2}\overline{\psi}\gamma^\nu\tau^3\psi \tag{14}$$

The Maxwell equations are of the form

$$-\partial_\mu F^{\mu\nu} = -e\,J^\nu$$

with the electric current given by

$$J^\nu = \frac{1}{2}\overline{\psi}\gamma^\nu Q_e\psi. \tag{15}$$

The timelike components of the baryon and isospin currants are the baryon density $\rho_B = J^0_B$ and the isospin density $\rho_a = J^0_a$. The conserved baryon and isospin charges are given by the following relations:

$$Q_B = \int d^3x\psi^+\psi = \int d^3x(\psi^+_n\psi_n + \psi^+_p\psi_p),$$

$$Q_3 = \frac{1}{2}\int d^3x\psi^+\sigma^3\psi = \frac{1}{2}\int d^3x(\psi^+_n\psi_n - \psi^+_p\psi_p),$$

The electric charge is defined by

$$Q = \int d^3x\psi^+Q_e\psi = \int d^3x\psi^+_p\psi_p \quad ,Q_e = \frac{1}{2}(I - \tau^3) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

The Dirac equations for nucleons that are obtained from the Lagrangian function have the following form

$$i\gamma^\mu D_\mu\psi - (M - g_{N\sigma}\varphi)\psi = 0. \tag{16}$$

The physical system is defined by the thermodynamic potential [16]

$$\Omega = -kTlnTr(e^{-\beta(H-\mu Q_B-\mu_3 Q_3)}) \tag{17}$$

where H stands for the Hamiltonian and is defined as

$$H = \sum_A\int d^3x\{\partial_0\Phi_A\pi^A - \mathcal{L}\}. \tag{18}$$

$\pi^A = \frac{\partial \mathcal{L}}{\partial(\partial_0 \Phi_A)}$ are momenta connected with fields $\Phi_A = \{\varphi,\ \omega_\mu,\ b_\mu, \psi\}$ which denote all fields considered in this model. The averaged charges can be obtained from the thermodynamic potential with the use of the relations

$$\frac{\partial \Omega}{\partial \mu} = - < Q_B >, \quad \frac{\partial \Omega}{\partial \mu_3} = - < Q_3 > . \tag{19}$$

In this paper the variational method based on the Feynman-Bogolubov inequality [17] is incorporated (see more details in [18])

$$\Omega \leq \Omega_1 = \Omega_0(m_{eff}) + < H - H_0 >_0 . \tag{20}$$

Ω_0 is the thermodynamic potential of the trial system which as effectively free quasiparticle system is described by the Lagrange function

$$\begin{aligned} \mathcal{L}_0(m_{eff}) = & \frac{1}{2}\partial_\mu \overline{\varphi} \partial^\mu \overline{\varphi} - \frac{1}{2} m^2_{eff,\sigma} \overline{\varphi}^2 - \frac{1}{4} \overline{\Omega}_{\mu\nu} \overline{\Omega}^{\mu\nu} + \frac{1}{2} m^2_{eff,\omega} \overline{\omega}_\mu \overline{\omega}^\mu \\ & - \frac{1}{4} \overline{G}^a_{\mu\nu} \overline{G}^{a\mu\nu} + \frac{1}{2} \sum_a m^2_{eff,\rho,a} \overline{b}^a_\mu \overline{b}^{a\mu} \\ & + \overline{\psi}(i\gamma^\mu \overline{D}_\mu - m_{eff,N})\psi \end{aligned} \tag{21}$$

with the field tensor now defined by

$$\overline{G}^a_{\mu\nu} = \partial_\mu \overline{b}^a_\nu - \partial_\nu \overline{b}^a_\mu$$

and

$$\overline{\Omega}_{\mu\nu} = \partial_\mu \overline{\omega}_\nu - \partial_\nu \overline{\omega}_\mu .$$

The fields Φ_A can be written as a sum of two components, the effectively free quasiparticle fields $\overline{\Phi}_A = \overline{\varphi}, \overline{\omega}_\mu, \overline{b}^a_\mu$ and the classical boson condensates $\xi_A = \sigma, w, b$

$$\Phi_A = \overline{\Phi}_A + \xi_A, \tag{22}$$

which in the case of particular meson fields results in

$$\varphi = \overline{\varphi} + \sigma \tag{23}$$

$$\omega_\mu = \overline{\omega}_\mu + w_\mu,\ \ w_\mu = \delta_{\mu,0} w \tag{24}$$

$$\rho^a_\mu = \overline{b}^a_\mu + \beta^a_\mu,\ \ \beta^a_\mu = \delta^{a,3}\delta_{\mu,0}\, b \tag{25}$$

Both the boson and fermion masses (m_B, m_F) as well as the $\xi_A = \{\sigma,\ w,\ b\}$ fields are treated as the variational parameters in the effective potential. The covariant derivative for the trial system is

$$\overline{D}_\mu = \partial_\mu + \frac{1}{2} i g_\rho \beta^a_\mu \tau^a + i g_\omega w_\mu \tag{26}$$

This introduces the fermion interaction with homogenous boson condensate w_μ, β^a_μ. The fermion quasiparticle obeys the Dirac equation

$$(i\gamma^\mu \overline{D}_\mu - m_{eff,N})\psi = 0 \tag{27}$$

Figure 1.



Nucleon effective mass $m_{eff,N} = M - g_{N\sigma}\sigma$ as the function of the baryon density n_B (fm^{-3}).

The masses entering the Lagrangian function $L_0(m_{eff})$ of the trial system calculated from the extremum conditions

$$\frac{\partial\Omega_1}{\partial m^2_{eff,\sigma}} = 0, \quad \frac{\partial\Omega_1}{\partial m^2_{eff,\omega}} = 0, \quad \frac{\partial\Omega_1}{\partial m^2_{eff,\rho}} = 0, \quad \frac{\partial\Omega_1}{\partial m_{eff,N}} = 0$$

now are the effective once. They are given by the following equations

$$\begin{aligned} m^2_{eff,\sigma} &= m_s^2 + 3g_3 < \overline{\varphi}^2 >_0 + 2b_2\sigma + 3b_3\sigma^2 \\ &- g^2_{\omega\sigma}(w^2 + < \overline{\omega}_\mu\overline{\omega}^\mu >_0) - g^2_{\rho\sigma}(b^2 + < \overline{b}^a_\mu\overline{b}^{a\mu} >_0), \end{aligned} \tag{28}$$

$$\begin{aligned} m^2_{eff,\omega} &= (M_\omega - g_{\sigma\omega}\sigma)^2 + g^2_{\omega\sigma} < \overline{\varphi}^2 >_0 + c_3(w^2 + < \overline{\omega}_\mu\overline{\omega}^\mu >_0) \\ &+ 2\Lambda_v(g_\rho g_\omega)^2(b^2 + < \overline{b}^a_\mu\overline{b}^{a\mu} >_0) \end{aligned} \tag{29}$$

$$\begin{aligned} m^2_{eff,\rho,a} &= (M_\rho - g_{\rho\sigma}\sigma)^2 + g^2_{\rho\sigma} < \overline{\varphi}^2 >_0 + 2(g_\rho g_\omega)^2\Lambda_v(w^2 + < \overline{\omega}_\mu\overline{\omega}^\mu >_0) \\ &- g_\rho^2 b^2(1-\delta_{a,3}) - \frac{3}{2}g_\rho^2\sum_{c\neq a} < \overline{b}^c_\mu\overline{b}^{c\mu} >_0^2 \\ &\xrightarrow[T=0]{} (M_\rho - g_{\rho\sigma}\sigma)^2 - g_\rho^2 b^2(1-\delta_{a,3}) + 2(g_\rho g_\omega)^2\Lambda_v w^2 \end{aligned} \tag{30}$$

$$m_{eff,N} = M - g_{N\sigma}\sigma. \tag{31}$$

In the presence of the ρ field condensation b the ρ mass $m_{eff,\rho,a}$ is splitting. This is a result of the 'nonabelian' character of the $SU(2)$ symmetry which is explicitely broken by the M_ρ term. In the homogeneous case the free energy reaches the minimum at σ. The nucleon effective mass $m_{eff,N}$ dependence on the baryon density n_B is presented in Fig. 1. Results are shown for the TM1 and NL3 parameter sets without additional coupling and for NL3 supplemented by the vector-scalar coupling.

The meson effective masses as functions of the baryon density n_B are presented in Figs. 2 and 3. This figure provides a comparison of the meson effective masses obtained



Figure 2. The mesons ω and ρ effective mass as the function of the baryon density n_B (fm^{-3}).



Figure 3. The mesons ω and σ effective mass as the function of the baryon density n_B (fm^{-3}).

for different values of the coupling constants. Straight dotted lines show the ω and ρ masses in the simplest model with no additional nonlinear terms. When this model is enlarged by the nonlinear vector-isoscalar and vector-isovector and the vector-scalar interactions the effective masses are obtained. As baryon number increases the ω meson effective mass increases and then become smaller. The ρ meson effective mass increases with the increasing baryon number.

The symmetry energy density (for $T = 0$) $\varepsilon_s = E_s/n_B$ can be evaluated taking into account the contributions of the ρ meson field to the energy of the system

$$E_s = \frac{1}{2}m_{eff,\rho}^2 b^2 = \frac{1}{2}(M_\rho - g_{\rho\sigma}\sigma)^2 b^2 + \Lambda_v(g_\rho g_\omega)^2 w^2 b^2 \qquad (32)$$



Figure 4. The symmetry energy E_s as the function of the baryon density n_B (fm^{-3}).



Figure 5. The static homogeneous solution for the ρ meson field as the function of the baryon density n_B (fm^{-3}).

Using the equation of motion the static homogeneous solution for the ρ field takes the following form

$$b = \frac{g_\rho}{m^2_{eff,\rho}} Q_3 = \frac{g_\rho n_B}{2m^2_{eff,\rho}} \delta \tag{33}$$

$$\delta = (n_n - n_p)/n_B \tag{34}$$

and is presented in Fig. 5. Taking into account the form of the equations (33) and (34) the symmetry energy now is given by

$$\varepsilon_s = \frac{g^2_\rho n_B}{8m^2_{eff,\rho}} \delta^2 \quad \rightarrow \quad \frac{g^2_\rho n_B}{8M^2_\rho} \delta^2 .$$

Parameter	NL3	TM1	QMC
E_0 (MeV)	-15.92	-16.26	-16.65
$k_{F,0}$ (fm^{-1})	1.24	1.29	1.297
ρ_0 (fm^{-3})	0.14	0.145	0.147
K (MeV)	170.24	281.53	284.71

Table 1. The nucleon symmetric phase properties (the binding energy E_0, the Fermi momentum, the density and the compresibility K) at the saturation point.

In the case of no additional coupling constants the result reproduces that of the simplest model without any nonlinearities. Including the $\omega-\rho$ and the vector-scalar coupling the ε_s is no longer linear with baryon density n_B. Including the $\omega-\rho$ and the vector-scalar coupling the ε_s is no longer linear with baryon number density n_B. For the purpose of the present study, we choose two basic RMF parameterizations (Table 3): NL3 [22], TM1 [23] and one based on QMC approach [14]. All parameters are chosen in such a form to reproduce the nucleon symmetric phase properties (the binding energy E_0, the Fermi momentum, the density and the compresibility K) at the satutation point (Table 1).

The case of static nonhomogeneous nuclear matter configuration can be described by the Euler-Lagrange equations (10, 12, 11, 16) taking the quantum average of these equations with respect to the effective quasiparticle system H_0 which in turn is obtained with the use of the effective quasiparticle Lagrangian $\mathcal{L}_0(m_{eff})$. The results for a spherically symmetric system become

$$\Delta\sigma(r) = \frac{\partial U_{eff}}{\partial\sigma}. \tag{35}$$

$$\Delta w(r) = m^2_{eff,\omega}\, w(r) - g_\omega\, Q_B, \tag{36}$$

$$\Delta b(r) = m^2_{eff,\rho}\, b(r) - g_\rho\, Q_3 \tag{37}$$

where the effective potential for the scalar meson is given by

$$\begin{aligned} U_{eff} = {} & (g_2 < \overline{\varphi}^2 >_0 + g_{\omega\sigma} M_\omega (w^2 + < \overline{\omega}_\mu \overline{\omega}^\mu >_0) + g_{\rho\sigma} M_\rho (b^2 + < \overline{b}^a_\mu \overline{b}^{a\mu} >_0))\sigma \\ & + \frac{1}{2}(m_s^2 + 3g_3 < \overline{\varphi}^2 >_0 - g^2_{\omega\sigma}(w^2 + < \overline{\omega}_\mu \overline{\omega}^\mu >_0) - g^2_{\rho\sigma}(b^2 + < \overline{b}^a_\mu \overline{b}^{a\mu} >_0))\,\sigma^2 \\ & + \frac{1}{3} g_2 \sigma^3 + \frac{1}{4} g_3 \sigma^4. \end{aligned} \tag{38}$$

The construction of U_{eff} is such that

$$m^2_{eff,\sigma} = \frac{\partial^2 U_{eff}}{\partial\sigma^2}.$$

The ρ condensate has a mass ($m_{eff,\rho} = m_{eff,\rho,3}$ at $T=0$)

$$\begin{aligned} m^2_{eff,\rho} = {} & (M_\rho - g_{\rho\sigma}\sigma)^2 + g^2_{\rho\sigma} < \overline{\varphi}^2 >_0 + 2(g_\rho g_\omega)^2 \Lambda_v (w^2 + < \overline{\omega}_\mu \overline{\omega}^\mu >_0) \\ & - g_\rho^2 \sum_{c\neq 3} < \overline{b}^c_\mu \overline{b}^{c\mu} >_0^2 . \end{aligned} \tag{39}$$



Figure 6. Profile of the ω meson field inside the ^{208}Pb nucleus ($g_{\omega\sigma} = g_{\rho\sigma} = 0$ case).

In the obtained results the thermal contributions from meson fields are present, however, they are not involved in calculations since meson masses are too large to be relevant. The Dirac equation for nucleons (i=p,n) has the following form

$$\{-i\alpha\nabla + V(r) + \beta[M - S(r)]\}\psi_i = \epsilon_i\psi_i \tag{40}$$

with the potentials:

$$V(r) = g_\omega\, w(r) + g_\rho\, \sigma^3 b(r) + eQ_e A_0(r),$$

$$S(r) = g_{N\sigma}\sigma(r).$$

In the non-honomegenous nuclear matter distribution the effective masses are space dependent and act rather like external potentials. The self-consistent equations (35, 36, 37, 40), namely the Dirac equation with potential terms for the nucleons and the Klein-Gordon type equations with sources for the mesons and the photon are the base for the numerical description of nuclei [19]. The simplified solutions (cf. Fig. 6 and Fig.4) can be obtained assuming the Fermi shape for the neutron and proton density distribution [7]. Spherically symmetric solutions for the ω and ρ meson fields for the ^{208}Pb nucleus ($g_{\omega\sigma} = g_{\rho\sigma} = 0$ case) are presented in Fig. 6 and Fig. 7. More realistic calculations, which will be the subject of future investigations, can be obtained with the use of the self-consistent Thomas-Fermi approximation [20], [21]. In the case of the ω meson the influence of the $\omega - \rho$ coupling is rather small whereas there exists a marked difference in the ρ meson field profile when the $\omega - \rho$ coupling is present (cf. (Fig. 7)).

3. Conclusions

The model presented here includes the original Walecka model with the nucleon fields and the kinetic and mass terms for scalar and vector mesons. Scalar nonlinear terms



Figure 7. Profile of the ρ meson field inside the ^{208}Pb nucleus ($g_{\omega\sigma} = g_{\rho\sigma} = 0$ case).

g_ρ[7]	Λ_v[7]	$R_p\,(fm)$[7]	$(R_n - R_p)\,(fm)$[7]	$m_{eff,\omega}\,(MeV)$	$m_{eff,\rho}\,(MeV)$
8.92	0	5.460	0.280	782.5	763.0
10.75	0.025	5.469	0.209	785.4	1141.6

Table 2. The meson effective masses inside the ^{208}Pb nucleus ($g_{\omega\sigma} = g_{\rho\sigma} = 0$ case).

were proposed by Boguta and Bodmer. Boguta and Price introduced quartic vector self-interaction. The chosen form of the Lagrange function which was supplemented by the additional omega-rho coupling was inspired by the work by Horowitz and Piekarewicz [7] and by the QMC model [14]. This additional nonlinear meson interaction can be explained in terms of effective meson masses that are modified in the nuclear medium. The additional nonlinear couplings Λ_V and $g_{\omega\sigma}$ and $g_{\rho\sigma}$ change the density dependence of the omega and rho fields, whereas Λ_V itself influences the density dependence of the symmetry energy. Taking into account these additional interaction terms the selfconsistent equations for vector mesons were obtained. The nucleon and scalar self-interaction generates different from zero expectation value of the scalar field and the effective nucleon mass $m_{eff,N}$. The mutual interactions of vector meson fields give as a result the effective meson masses $m_{eff,\omega}$ and $m_{eff,\rho}$. The appearance of the effective meson masses changes the energy symmetry of the system and has an influence on the behaviour of the ρ field. All calculations were performed with the use of NL3 [22] and TM1 [23] parameter sets. They not only correctly reproduce nuclear properties but also are consistent with the idea of naturalness as well. The force of NL3 stems from the fit including exotic nuclei, neutron radii, and information on giant resonances. The parameter set named TM1 has been applied to deformed nuclei, triaxal deformations, nuclear excitations [10, 24, 25]. The last column mark by QMC represents parameters produced by the Quark Meson Coupling model (QMC) [9, 14].

Parameter	NL3 [22]	NL3 [7]	NL3 [7]	TM1 [23]	QMC [14]
M	$938\,MeV$	$938\,MeV$	$938\,MeV$	$938\,MeV$	$939\,MeV$
M_w	$795.4\,MeV$	$795.4\,MeV$	$795.4\,MeV$	$783\,MeV$	$783\,MeV$
M_ρ	$763\,MeV$	$763\,MeV$	$763\,MeV$	$770\,MeV$	$770\,MeV$
m_s	$492\,MeV$	$492\,MeV$	$492\,MeV$	$511.2\,MeV$	$550\,MeV$
$g_2 = \kappa/2$	$12.17\,fm^{-1}$	$12.17\,fm^{-1}$	$12.17\,fm^{-1}$	$7.23\,fm^{-1}$	$9.62\,fm^{-1}$
$g_3 = \lambda/6$	-36.259	-36.259	-36.259	0.6183	61.24
$g_{N\sigma}{=}g_s$	10.138	10.138	10.138	10.0289	6.9667
g_ω	13.285	13.285	13.285	12.6139	5.8561
g_ρ	8.941	9.214	10.751	9.2644	8.3803
c_3	0	0	0	71.3075	0
Λ_v	0	0.005	0.025	0	0
$g_{\omega\sigma}$	0	$2/3\,g_{N\sigma}$	$2/3\,g_{N\sigma}$	0	0.6617
$g_{\rho\sigma}$	0	$2/3\,g_{N\sigma}$	$2/3\,g_{N\sigma}$	0	0.6617

Table 3. Parameter sets for the Lagrangian (1).

References

[1] Boguta J. and Bodmer A.R. 1977 *Nucl. Phys.* **A292** 413

[2] Serot B.D., Walecka J.D. 1997 *Recent Progress in Quantum Hadrodynamics Int. J. Mod. Phys.* **E6** 515

[3] Ring P 1996 *Prog. Part. Nucl. Phys.* **37** 193

[4] Glendenning N.K. 1987 *Z. Phys., A327* 295; Also see in *Compact Stars* by N. K Glendenning Sringer-Verlag, New York, 1997

[5] Weber F.*Pulsars as Astrophysical Laboratories for Nuclear and Particle Physics,* IOP Publishing, Philadelphia, 1999

[6] Gmuca S.J. 1991 *J.Phys.* **G17** 1115

[7] Horowitz C. J. and Piekarewicz J. 2001 *Phys. Rev. Lett.* **86**, 5647

[8] Li G.Q., Ko C.M., Brown G.E. 1996 *Nucl. Phys.***A606**, 568

[9] Guichon P.A.M., *Phys. Lett.* **B 200** 235; Guichon P.A.M., Saito K., Rodionov E., Thomas A.W. *Nucl. Phys.* **A 601** 394; Saito K., Tsushima K., Thomas A.W. *Phys. Rev.* **C 55** 2637;

[10] Hirata D, Toki H. and Tannihata I. 1995 *Nucl. Phys.* **A589** 239

[11] Serot B.D. and Walecka J.D. 1986 *Adv. Nucl. Phys.* **16** 1

[12] Serot B.D. 1992 *Rep. Prog. Phys.* **55** 1855

[13] Bodmer A.R. 1991 *Nucl. Phys.* **A526** 703

[14] Saito K., Tsushima K., Thomas A.W. *Phys. Lett.* **B 406** 287;

[15] Horowitz C. J., Pollock S. J., Souder P. A. and Michaels R. *Preprint* nucl-th/9912038

[16] Fetter A.L. and Walecka J.D. *Quantum Theory of Many-Particle Systems,* McGraw-Hill, New York 1971.

[17] Feynman R., *Statistical Mechanics,* W.A.Benjamin, Inc. Reading 1972.
Mańka R., Kuczyński J. and Vitiello G.V. 1986 *Nucl. Phys.* **B276** 533; Mańka R. and Vittiello G.V. 1990 *Annals of Physics* **199** 61

[18] Mańka R., Bednarek I. and Przybyła 2000 *Phys. Rev.* **C62** 015802

[19] Ring P., Gambhir Y. K. and Lalazissis G. A. 1997 *Comput. Phys. Commun.* **105** 77

[20] Menezes D.P.,Providencia C. 1999 *Nucl. Phys.* **A 650**, 283

[21] Menezes D.P.,Providencia C. 1999 *Phys. Rev.* **C 60**, 024313

[22] Lalazissis G.A. , König J. and Ring P. 1997 *Phys. Rev.* **C55**, 540;.P.-G.Reinhard, Rufa M. , Maruhn J., Greiner W. and Friedrich J. 1986 *Z. Phys. A - Atomic Nuclei* **323** 13
[23] Sugahara Y. and Toki H. 1994 *Prog. Theo. Phys* **92** 803
[24] Hirata D, Sumiyoshi K. Tanihata I., Tachibana Y. and Toki H. 1997 *Nucl. Phys.* **A616** 438c
[25] Ma Z.Y., Toki H., Chen B. Q. and Van Giai N. 1997 *Prog.Theor. Phys.* **98** 917